



An Australian Government Initiative

Dementia Risk Reduction: A Practical Guide for Health and Lifestyle Professionals



March 2010



CONTENTS

- ACKNOWLEDGEMENTS AND DISCLAIMERS iv
- 1. INTRODUCTION 1
 - 1.1 About this guide 1
 - 1.2 What are the risk factors for dementia? 1
 - 1.3 Why is dementia risk reduction important?..... 2
 - 1.4 References 3
- 2. SUMMARY OF EVIDENCE AND RECOMMENDATIONS 4
 - 2.1 Chronic disease risk factors and dementia 4
 - 2.2 Vascular disease risk factors and dementia..... 4
 - 2.3 Psychosocial factors and dementia 4
 - 2.4 Other risk factors and dementia..... 5
- 3. ALCOHOL 6
 - 3.1 Moderate alcohol may reduce dementia risk, but heavy drinking increases risk.... 6
 - 3.2 Alcohol and dementia risk – the evidence..... 6
 - 3.3 Promoting healthy use of alcohol in professional practice..... 6
 - 3.4 Alcohol resources for clients 7
 - 3.5 References 7
- 4. BLOOD PRESSURE..... 8
 - 4.1 High blood pressure increases dementia risk..... 8
 - 4.2 Blood pressure and dementia risk – the evidence 8
 - 4.3 How does blood pressure impact on dementia risk? 8
 - 4.4 Blood pressure resources for clients 8
 - 4.5 References 9
- 5. BODY WEIGHT 10
 - 5.1 Obesity increases dementia risk..... 10
 - 5.2 Body weight and dementia risk – the evidence..... 10
 - 5.3 How does excess weight impact on dementia risk? 10
 - 5.4 Body weight assessment and management in professional practice..... 10
 - 5.5 Weight management resources for clients 11
 - 5.6 References 11
- 6. CHOLESTEROL 12
 - 6.1 High cholesterol in midlife increases dementia risk 12
 - 6.2 Cholesterol and dementia risk – the evidence 12
 - 6.3 How does cholesterol impact on dementia risk?..... 12
 - 6.4 Cholesterol resources for clients 12
 - 6.5 References 13

7. DEPRESSION	14
7.1 A history of depression is associated with increased dementia risk	14
7.2 Depression and dementia – the evidence	14
7.3 Depression resources for professionals	14
7.4 Depression resources for clients.....	15
7.5 References	15
8. DIABETES	16
8.1 Diabetes increases dementia risk.....	16
8.2 Diabetes and dementia risk – the evidence.....	16
8.3 Diabetes resources for professionals.....	16
8.4 Diabetes resources for patients.....	17
8.5 References	17
9. DIET.....	18
9.1 A brain healthy diet may reduce dementia risk.....	18
9.2 Diet and dementia risk – the evidence	18
9.3 Promoting healthy diets in professional practice	19
9.4 Healthy diet resources for clients	19
9.5 References	20
10. HEAD INJURY.....	21
10.1 Head injury increases the risk of dementia	21
10.2 Head injury and dementia – the evidence.....	21
10.3 Prevention of head injury in professional practice	21
10.4 Injury prevention information for clients.....	22
10.5 References.....	22
11. MENTAL STIMULATION	23
11.1 Mental stimulation may reduce dementia risk.....	23
11.2 Mental activity and dementia risk – the evidence.....	23
11.3 Promoting mental activity in professional practice.....	23
11.4 Mental activity resources for clients	24
11.5 References.....	25
12. PHYSICAL ACTIVITY.....	26
12.1 Physical exercise may reduce dementia risk.....	26
12.2 Physical Activity and dementia risk – the evidence.....	26
12.3 Promoting physical activity in professional practice	26
12.4 Physical activity resources for clients	27
12.5 References.....	27

13. SMOKING	28
13.1 Smoking increases dementia risk	28
13.2 Smoking and dementia risk – the evidence	28
13.3 Promoting smoking cessation in professional practice	28
13.4 Smoking cessation resources for clients	29
13.5 References.....	29
14. SOCIAL ACTIVITY	30
14.1 Social interaction may reduce dementia risk	30
14.2 Social Activity and dementia risk – the evidence.....	30
14.3 How does social activity reduce dementia risk?	30
14.4 Promoting social activity in professional practice.....	30
14.5 Social activity resources for clients	31
14.6 References.....	31
APPENDIX.....	32
Dementia risk reduction summary sheet (one page) for distribution to clients.....	32

ACKNOWLEDGEMENTS AND DISCLAIMERS

Dementia Risk Reduction: A Practical Guide for Health and Lifestyle Professionals was written by Dr Maree Farrow, Research Fellow, Alzheimer's Australia Vic, and was prepared as part of Alzheimer's Australia Vic's involvement in the Dementia Collaborative Research Centre: Early Diagnosis and Prevention.

The [Dementia Collaborative Research Centres](#) are an Australian Government funded initiative established to advance Australian research into dementia and the translation of research into clinical practice. The three Centres each focus on a different area of dementia research:

- Assessment and better care
- Early diagnosis and prevention
- Carers and consumers

Many thanks to Associate Professor Michael Woodward and Dr David Lie for helpful suggestions.

Whilst appreciable care has been taken in the preparation of this guide, Alzheimer's Australia and its member organisations accept no responsibility for any inaccuracies, omissions or information that may be perceived as misleading. The guide is for information purposes only, and is designed as a general reference and link to further information about preventive health. Alzheimer's Australia is not responsible for the results of any actions taken by any person on the basis of any information in this publication.

Copyright in the product sample templates, Commonwealth logo, photographs and graphic layouts reproduced from the *Dementia Research Graphic Design Standards Manual* is owned by or licensed to the Commonwealth of Australia and published with the permission of the Commonwealth of Australia on the condition reproduction occurs for non-commercial use and promotes or benefits selected Commonwealth approved dementia initiatives and programs. All commercial and other rights are reserved.

Important notice: this work may not be a Commonwealth publication or product

The views expressed in this work are the views of its author(s) and not necessarily those of the Commonwealth of Australia. Despite any permitted use of the *Dementia Research Graphic Design Standards Manual* copyright or licensed material, the reader needs to be aware that the information contained in this work is not necessarily endorsed, and its contents may not have been approved or reviewed, by the Australian Government Department of Health and Ageing.

ISBN 978-1-921570-12-4

© Alzheimer's Australia 2010
ABN 79 625 582 771 ARBN 066 779 557

1. INTRODUCTION

1.1 About this guide

This guide provides information for health and lifestyle professionals about modifiable risk and protective factors for dementia. Each section explains the evidence for the association of that factor with dementia risk and provides a practical guide to the resources available to health and lifestyle professionals to assist them to work with their clients to address factors of concern. Where possible, hyperlinks are provided to those resources.

The guide aims to provide health and lifestyle professionals with a quick reference source to enable them to offer strategies to clients for maintaining their cognitive health and reducing their dementia risk. As many of the risk factors for dementia overlap with those for other chronic conditions that can be addressed by preventative health, the guide draws on existing guidelines and resources, and provides links to these.

Alzheimer's Australia's Mind your Mind is a community education program designed to inform people about what they can do to reduce their risk of dementia. It is based on the scientific evidence that is described in this guide. The program consists of seven 'signposts' toward dementia risk reduction:

- ◆ Mind your Body – be physically active
- ◆ Mind your Brain – keep your brain active
- ◆ Mind your Diet – eat healthily
- ◆ Mind your Habits – don't smoke and drink alcohol only in moderation
- ◆ Mind your Head – protect your head from injury
- ◆ Mind your Health Checks – check and manage blood pressure, cholesterol, blood sugar and weight
- ◆ Mind your Social Life – participate in social activities

The risk and protective factors associated with each of the Mind your Mind signposts and resources for addressing these are covered in this guide.

1.2 What are the risk factors for dementia?

Non-modifiable risk factors

Age is the most important risk factor for dementia, with prevalence increasing exponentially after age 65. In 2009, the prevalence of dementia in Australia was estimated to rise from 1.7% of men and 1.3% of women aged 65-69, to 37% of men and 47% of women aged 95 and over [1].

Autosomal dominant gene mutations have been identified for some rare causes of dementia. These account for only a small percentage of dementia cases, and usually result in younger onset dementia (diagnosed before 65 years of age). For example, variations in specific genes are known to cause familial Alzheimer's disease, forms of frontotemporal dementia and Huntington's disease [2].

People with Down syndrome are at very high risk of developing Alzheimer's disease [2]. A susceptibility gene, the apolipoprotein E – epsilon 4 allele, is associated with a higher risk of developing Alzheimer's disease and vascular dementia [2]. Routine testing for this gene is not currently recommended.

Having a family history of dementia is associated with an increased risk. Those with a first degree relative having Alzheimer's disease have a 2-3 fold increased risk of developing the disease themselves [3].

The prevalence of Alzheimer's disease is higher in females than males [1]. The prevalence of vascular dementia is higher in males [1].

Modifiable risk factors

Because of the influence of non-modifiable risk factors, especially age and genetics, dementia cannot be definitely prevented, but there is growing evidence that addressing modifiable risk factors can reduce the risk or delay the onset of dementia.

Epidemiological research shows that certain medical and lifestyle factors are consistently associated with a higher or lower risk of developing dementia. Regular physical, mental and social activity are associated with reduced dementia risk, while vascular risk factors are associated with increased dementia risk [4]. Encouraging active lifestyles and effective treatment of vascular risk factors therefore has the potential to reduce the risk of dementia among the population.

Modifiable factors associated with dementia risk include:

- ◆ Alcohol consumption
- ◆ Blood pressure
- ◆ Body weight
- ◆ Cholesterol
- ◆ Depression
- ◆ Diabetes
- ◆ Diet
- ◆ Head injury
- ◆ Mental activity
- ◆ Physical activity
- ◆ Smoking
- ◆ Social activity

1.1 Why is dementia risk reduction important?

Many people, especially those with a family history of dementia, are interested in what they can do to prevent developing the illness themselves. Health and lifestyle professionals can assist clients by providing advice that several lifestyle and health factors are consistently associated with the risk of developing dementia. They can use recommended interventions to reduce the impact of modifiable risk factors for clients.

The incidence of dementia is growing rapidly with the ageing of the population. In Australia in 2010 there are an estimated 257,000 people with dementia. This is expected to increase to 1,131,000 by 2050 [1].

While there is no cure for dementia, addressing modifiable risk factors remains an important strategy for reducing the incidence of dementia. Modelling suggests that if the average onset age could be delayed by 5 years, incidence would be halved by 2050 [5].

Half of Australian adults remain unaware of the potential for dementia risk reduction. Some associate mental activity with reduced risk, but very few acknowledge the important link between vascular risk factors and dementia [6]. Health and lifestyle professionals can help address this by educating clients about the risk factors, helping clients make appropriate lifestyle changes, and referring them to their doctor to address medical risk factors.

Preventative health strategies aimed at maintaining cognitive health have the potential to have a significant impact on future numbers of people developing dementia, and health and lifestyle professionals have an important role to play in this. Dieticians, exercise professionals, pharmacists, psychologists, occupational therapists, practice nurses, community health workers and others who work with clients to improve their health can be involved in helping them maintain their cognitive health and reduce their risk of dementia.

1.4 References

1. Access Economics. *Keeping dementia front of mind: incidence and prevalence 2009-2050*. 2009, Alzheimer's Australia, Canberra.
2. Hsiung GY & Sadvnick AD. Genetics and dementia: risk factors, diagnosis, and management. *Alzheimers Dement*, 2007, 3:418-427.
3. Jarvik LF & Blazer D. Children of Alzheimer patients: an overview. *J Geriatric Psychiatry Neurology*, 2005, 18:181-186.
4. Woodward M, et al. *Dementia risk reduction: the evidence*. 2007, Alzheimer's Australia, Canberra.
5. Access Economics. *Delaying the onset of Alzheimer's disease: projections and issues*. 2004, Alzheimer's Australia, Canberra.
6. Farrow M. *Dementia risk reduction: what do Australians know?* 2008, Alzheimer's Australia, Canberra.

2. SUMMARY OF EVIDENCE AND RECOMMENDATIONS

2.1 Chronic disease risk factors and dementia

- ◆ Consumption of moderate amounts of alcohol has been associated with better cognitive function and reduced risk of dementia. However, excessive amounts over time can increase the risk.
- ◆ Obesity at midlife is associated with an increased risk of developing dementia, including Alzheimer's disease. In old age, those who are underweight or are losing weight are found to be at increased risk of dementia.
- ◆ Diets low in saturated fat and high in antioxidant rich fruits and vegetables have been associated with better brain function and reduced risk of dementia. What is good for the heart seems to be also good for the brain.
- ◆ Regular physical exercise in midlife and late-life is associated with better brain function and reduced risk of cognitive decline and dementia. Even simple exercise like walking has been shown to be beneficial.
- ◆ Smoking is associated with an increased risk of dementia. Former smokers have lower dementia risk, demonstrating the benefits of quitting.

The behavioural risk factors above are common among the Australian population. They contribute significantly to the burden of disease, due to their influence on chronic diseases such as diabetes, cardiovascular disease, respiratory disease and some cancers. They are also associated with the risk of dementia. Interventions by health and lifestyle professionals can help clients make lifestyle changes and reduce their risk of chronic disease including dementia.

2.2 Vascular disease risk factors and dementia

- ◆ High blood pressure, especially at midlife, is consistently associated with an increased risk of developing dementia. Treatment of hypertension has been found to reduce the risk of cognitive decline and dementia in several studies.
- ◆ High serum total cholesterol, especially at midlife, is associated with an increased risk of developing dementia, especially Alzheimer's disease.
- ◆ Diabetes is associated with an increased risk of developing dementia, including Alzheimer's disease and vascular dementia.

The physiological risk factors above contribute to cardiovascular disease, which is thought to be largely preventable. They also increase the risk of dementia. Lowering blood pressure in patients with hypertension, reducing blood levels of total cholesterol, and maintaining good glycaemic control in patients with diabetes have been demonstrated to reduce stroke and heart attack. They may also reduce the incidence of dementia. Assessment and treatment of vascular risk factors must be undertaken by a doctor. Health and lifestyle professionals should encourage clients to talk to their doctor about this. For some people, reliance on lifestyle changes will not be enough.

2.3 Psychosocial factors and dementia

- ◆ People with depression or high depressive symptoms have, on average, a higher risk of developing dementia.
- ◆ Higher participation in mentally stimulating activities is associated with reduced risk of cognitive decline and dementia. Those with a history of higher education, mentally demanding occupations or participation in mentally challenging leisure activities are consistently found to have a lower risk of developing dementia.
- ◆ Being more socially active is associated with reduced risk of cognitive decline and dementia. This has been shown in many studies measuring social engagement in different ways.

Depression, social isolation and lack of quality social support are risk factors for coronary heart disease. They are also associated with increased risk of dementia. Health and lifestyle professionals can provide advice and resources to assist clients remain mentally and socially active. Clients at risk of depression should be referred to a doctor. Such interventions are likely to have benefits for patients' mental and cognitive health and may reduce their risk of dementia.

2.4 Other risk factors and dementia

- ◆ Head injury, with loss of consciousness, is associated with an increased risk of subsequent cognitive decline and dementia.

Prevention of head trauma is important for a number of reasons, including its potential effect on dementia risk. Health and lifestyle professionals can encourage clients to follow current safety advice and legislation to prevent head injury. This includes wearing helmets when appropriate and preventing falls in older adults.

This guide points health and lifestyle professionals to the resources and guidelines for preventative health interventions recommended to address the dementia risk factors summarised above.

3. ALCOHOL

3.1 Moderate alcohol may reduce dementia risk, but heavy drinking increases risk

Consumption of moderate amounts of alcohol has been associated with better cognitive function and reduced risk of dementia. However, excessive amounts over time can increase the risk. Limiting consumption to the recommended no more than 2 drinks a day may help reduce the risk of developing dementia.

This section describes the current evidence that suggests alcohol consumption can contribute to dementia risk and provides links to resources on healthy alcohol use for health and lifestyle professionals and resources for patients.

3.2 Alcohol and dementia risk – the evidence

Several studies have shown that light to moderate alcohol consumption is associated with a lower risk of any dementia, Alzheimer's disease and cognitive impairment. There is insufficient evidence to promote alcohol consumption to non-drinkers as a means of reducing dementia risk. However, there may be benefits for those currently using alcohol moderately.

In a large study of 8,000 people in Rotterdam, followed up for an average of 7 years, there was a 45% lower risk of dementia in those who consumed 1-3 alcoholic drinks per day, compared to non-drinkers [1].

Two meta-analyses found reduced risks of around 30 to 40% for those drinking moderate levels of alcohol [2,3]. Both found significant effects for any dementia and Alzheimer's disease. One also found a significant effect for vascular dementia [2].

However, excessive alcohol consumption may increase the risk of dementia. Very heavy drinking over time can cause alcohol-related dementia. While a meta-analysis failed to detect an association between heavy drinking and dementia risk [2], several studies have shown this association. For example, in a study of 554 twins in Finland, followed for 25 years, binge drinking at least monthly in midlife was associated with a more than 3 fold increase in the risk of dementia after the age of 65 years [4].

The upper limit of alcohol consumption that is safe for maintaining cognitive function in old age has not been determined. Alcohol's potential harmful effects for other medical conditions also need to be taken into consideration. The National Health and Medical Research Council's [Australian Guidelines to Reduce Health Risks from Drinking Alcohol](#) recommend limiting alcohol intake to no more than 2 drinks on any day to reduce the risk of long term harm [5].

The benefits of alcohol may be produced through its favourable effects on the cardiovascular system, although there may be other mechanisms. Reducing inflammation, increasing HDL cholesterol, increasing cerebral blood flow, increasing insulin sensitivity and antioxidant properties have all been implicated as contributing to alcohol's protective effect [2,3].

3.3 Promoting healthy use of alcohol in professional practice

Everyone, including those concerned about their dementia risk, should be encouraged to drink alcohol responsibly.

The National Health and Medical Research Council's [Australian Guidelines to Reduce Health Risks from Drinking Alcohol](#) outline recommended drinking levels for reducing the

risk of harm. They recommend all adults limit alcohol consumption to no more than 2 drinks on any day to reduce risk of long term harm, and no more than 4 drinks on any day to reduce risk of short term harm.

The [SNAP Guide](#) details recommendations and guidelines for assessing someone's alcohol consumption and assisting those needing to reduce their drinking. The SNAP guide was developed for general practitioners, but much of the information could be used by other professionals.

The [Australian Drug Foundation](#) provides a range of information, reports and resources on alcohol.

3.4 Alcohol resources for clients

The [Mind your Habits](#) section of the Alzheimer's Australia Mind your Mind website provides information for the general public on alcohol and its relationship to dementia risk.

The National Health and Medical Research Council's [Australian Guidelines to Reduce Health Risks from Drinking Alcohol](#) outline recommended drinking levels for reducing the risk of short term and long term harm.

The [Australian Drug Foundation](#) provides a range of information, fact sheets and resources on alcohol and standard drinks, and contacts for helplines providing counselling and referral.

3.5 References

1. Ruitenbergh A, et al. Alcohol consumption and risk of dementia: the Rotterdam study. *Lancet*, 2002, 359:281-286.
2. Anstey KJ, et al. Alcohol consumption as a risk factor for dementia and cognitive decline: meta-analysis of prospective studies. *Am J Geriatr Psychiatry*, 2009, 17:542-555.
3. Peters R, et al. Alcohol, dementia and cognitive decline in the elderly: a systematic review. *Age and Ageing*, 2008, 37:505-512.
4. Jarvenpaa T, et al. Binge drinking in midlife and dementia risk. *Epidemiology*, 2005, 16:766-771.
5. National Health and Medical Research Council. *Australian guidelines to reduce health risks from drinking alcohol*. 2009, Commonwealth of Australia, Canberra.

4. BLOOD PRESSURE

4.1 High blood pressure increases dementia risk

High blood pressure, especially at midlife, is consistently associated with an increased risk of developing dementia. Treatment of hypertension has been found to reduce the risk of cognitive decline and dementia in several studies.

This section describes the current evidence that suggests assessment and control of blood pressure are important for reducing dementia risk and provides links to resources for clients.

4.2 Blood pressure and dementia risk – the evidence

Hypertension is a risk factor for cerebrovascular disease and vascular dementia. Hypertension in midlife is also associated with an increased risk of Alzheimer's disease. Blood pressure may decline in the preclinical stage of Alzheimer's and low blood pressure in late-life may be associated with increased risk [1].

Treatment of hypertension in old age has been identified in several studies to reduce the risk of cognitive decline and dementia. In a 4 year study of 2,400 patients, treatment reduced the risk of dementia by 55% [2].

Studies assessing long term use of antihypertensives from midlife show a cumulative reduction in risk of dementia for each year of treatment [1,3]. A study that followed hypertensive men from midlife found that for each additional year of treatment there was a further reduction in the risk of dementia [3].

Those treated for more than 12 years had a 60% reduced risk of any dementia and 65% reduced risk of Alzheimer's disease compared to those never treated. The risk after 12 years of treatment was similar to those with normal blood pressure.

4.3 How does blood pressure impact on dementia risk?

Hypertension may contribute to both cerebrovascular pathology and Alzheimer's disease pathology [4]. Hypertension may lead to blood vessel endothelial dysfunction, mini strokes and cerebral ischaemia. Endothelial changes could also affect microvascular clearance of beta amyloid (the protein that accumulates into plaques) and exacerbate Alzheimer's disease pathology.

4.4 Blood pressure resources for clients

It is important for everyone, including those concerned about their risk of developing dementia, to know that hypertension is a risk factor for dementia that is easily treated. Hypertension often has no symptoms, so it is recommended that adults have their blood pressure measured by a doctor at least every 2 years. Those with other risk factors or conditions may require more regular checks.

Hypertension, if present, should be effectively treated under medical supervision. Untreated hypertension increases the risk of dementia, as well as heart disease, stroke and kidney disease. Treatment may involve lifestyle modification (diet and physical activity) and/or medication.

The [Mind your Health Checks](#) section of the Alzheimer's Australia Mind your Mind website provides information for the general public on blood pressure and its relationship to dementia risk.

The [Heart Foundation](#) has a range of information sheets on blood pressure, including self-measurement and the effects of dietary salt.

The [Stroke Foundation](#) provides information on blood pressure and runs an annual 'Know your Numbers' campaign to promote awareness of the importance of checking blood pressure.

4.5 References

1. Haag MD, et al. Duration of antihypertensive drug use and risk of dementia. A prospective cohort study. *Neurology*, 2009, 72:1727-1734.
2. Forette F, et al. The prevention of dementia with antihypertensive treatment. New Evidence from the Systolic Hypertension in Europe (Syst-Eur) study. *Arch Intern Med*, 2002, 162:2046-2052.
3. Peila R, et al. Reducing the risk of dementia. Efficacy of long-term treatment of hypertension. *Stroke*, 2006, 37:1165-1170.
4. Knopman DS. Hypertension and late-life dementia. A real link? *Neurology*, 2009, 72:1716-1717.

5. BODY WEIGHT

5.1 Obesity increases dementia risk

Obesity at midlife is associated with an increased risk of developing dementia, including Alzheimer's disease. Weight reduction has not as yet been shown to reduce the risk of dementia. In old age, those who are underweight or are losing weight are found to be at increased risk of dementia.

This section describes the current evidence that suggests maintaining a healthy weight is important for reducing dementia risk and provides links to resources for health and lifestyle professionals and for patients.

5.2 Body weight and dementia risk – the evidence

Obesity, particularly central obesity, at midlife is associated with increased risk of later developing dementia [1]. Whether midlife weight loss for those who are obese can reduce the risk of late life dementia has not yet been determined.

In early old age (65 to 75 years), there is a U-shaped relationship between BMI and dementia risk, with increased risk found for those who are underweight or obese [1].

In late life (after 75 years), lower BMI and weight loss have been associated with an increased risk of dementia [1]. Weight loss may precede dementia onset by more than 10 years.

A recent review summarised the findings of the complex relationship between weight and dementia across the lifespan as follows [1]:

- ◆ central adiposity in middle age predicts dementia in old age
- ◆ the relation between high adiposity and dementia is attenuated with older age
- ◆ waist circumference in old age, a measure of central adiposity, may be a better predictor of dementia than BMI
- ◆ lower BMI predicts dementia in the elderly
- ◆ weight loss may precede dementia diagnosis by decades

While there is still more to learn, the available evidence suggests that maintaining a healthy weight over the life course is a strategy likely to optimise both body and brain health. There is clinical trial evidence that weight loss lowers blood pressure, improves blood lipids and insulin resistance, and positively affects other factors associated with cardiovascular and dementia risk.

5.3 How does excess weight impact on dementia risk?

There are a number of potential mechanisms by which obesity may increase dementia risk [1]. Insulin resistance and hyperinsulemia can result from obesity and may play a role in reducing beta amyloid clearance from the brain, increasing the risk of Alzheimer's disease. Adipose tissue produces adipokines and cytokines, related to hyperinsulemia and inflammation that in turn have effects on the brain. Obesity is also associated with increased risk for cardiovascular and cerebrovascular disease, which increase the risk of dementia.

5.4 Body weight assessment and management in professional practice

People who are overweight or obese, including those concerned about their risk of developing dementia, should be educated about lifestyle choices to help maintain a healthy weight. Obesity in midlife increases the risk of dementia, as well as heart disease and a range of other conditions.

The [SNAP Guide](#) provides detailed information on assessing BMI and waist circumference and assisting patients who need to lose weight. The SNAP guide was developed for general practitioners, but much of the information could be used by other professionals.

The National Health and Medical Research Council's [Clinical practice guidelines for the management of overweight and obesity in adults](#) outline the aetiology of obesity and provide guidelines for assessing, measuring and treating overweight and obesity.

See also the sections in this guide on physical activity (section 11, p. 24) and healthy eating (section 8, p.16).

5.5 Weight management resources for clients

The [Mind your Health Checks](#) section of the Alzheimer's Australia Mind your Mind website provides information for the general public on body weight and its relationship to dementia risk.

The [Measure Up](#) campaign website provides information and resources and links to further information on weight, physical activity and healthy eating.

The Department of Health and Ageing [Healthy Weight](#) website provides information, tips and resources for maintaining a healthy weight.

The [Heart Foundation](#) provides information for consumers on weight, physical activity and healthy eating.

The [National Physical Activity Guidelines](#) from the Australian Government Department of Health and Ageing outline the minimum levels of physical activity required to gain a health benefit and ways to incorporate physical activity into everyday life.

The [Australian Dietary Guidelines](#) highlight the groups of foods and lifestyle patterns that promote good nutrition and health. Booklets and pamphlets summarising the guidelines for the general public are available at the website.

5.6 References

1. Luchsinger JA & Gustafson DR. Adiposity and Alzheimer's disease. *Current Opinion in Clinical Nutrition and Metabolic Care*, 2009, 12: 15-21.

6. CHOLESTEROL

6.1 High cholesterol in midlife increases dementia risk

High total cholesterol, especially at midlife, is associated with an increased risk of developing dementia, especially Alzheimer's disease. Treatment with statins (cholesterol lowering drugs) has been found to be associated with reduced risk of dementia in several studies, but there are some conflicting findings.

This section describes the current evidence that suggests assessment and control of cholesterol levels are important for reducing dementia risk and provides links to resources for clients.

6.2 Cholesterol and dementia risk – the evidence

Recent reviews have concluded that high midlife total serum cholesterol is associated with an increased risk of any dementia and of Alzheimer's disease [1,2]. While there are some studies that have not found this association, the majority of studies show that high midlife cholesterol increases the risk of later developing dementia by around 2 times.

A meta-analysis of studies measuring cholesterol at late life and following participants over a few years found no association between late life cholesterol and dementia risk [2]. Another study found that a decrease in cholesterol level from midlife to late life was associated with an increased risk of Alzheimer's disease [3]. Cholesterol levels may decline in the preclinical stages of Alzheimer's [1,2].

Several studies have shown reduced risk of dementia in those treated with statins, but a meta-analysis found only a small non-significant benefit [4].

6.3 How does cholesterol impact on dementia risk?

Some epidemiological and experimental studies suggest the cellular cholesterol level to be involved in the pathogenesis of AD [4]. High cholesterol levels may accelerate the production of beta amyloid [4]. Autopsy findings indicated that low midlife total cholesterol was associated with a lower number of plaques and tangles [1].

6.4 Cholesterol resources for clients

It is important for everyone, including those concerned about their risk of developing dementia, to know that high cholesterol is a risk factor for dementia that is easily treated. High cholesterol often has no symptoms, so it is recommended that cholesterol should be measured in adults 45 and over at least every 5 years, and those with other risk factors or conditions may require earlier and/or more regular checks.

High cholesterol, if present, should be effectively treated under medical supervision. Untreated it increases the risk of dementia, as well as heart disease and stroke. Treatment may involve lifestyle modification (diet and physical activity) and/or medication.

The [Mind your Health Checks](#) section of the Alzheimer's Australia Mind your Mind website provides information for the general public on cholesterol and its relationship to dementia risk.

The [Heart Foundation](#) has a range of information on cholesterol, how to lower cholesterol levels, and the right foods to eat.

The [Better Health Channel](#) provides a fact sheet on cholesterol for consumers.

The [ABC Health and Wellbeing](#) website has a fact file on cholesterol that explains what it is, how it is tested, and how to treat high cholesterol.

6.5 References

1. Kivipelto M & Solomon A. Cholesterol as a risk factor for Alzheimer's disease – epidemiological evidence. *Acta Neurol Scand*, 2006, 114(Suppl. 185):50-57.
2. Anstey KJ, et al. Cholesterol as a risk factor for dementia and cognitive decline: a systematic review of prospective studies with meta-analysis. *Am J Geriatr Psychiatry*, 2008, 16:343-354.
3. Stewart R, et al. Twenty-six-year change in total cholesterol levels and incident dementia: the Honolulu-Asia Aging Study. *Arch Neurol*, 2007, 64:103-107.
4. Zhou B, et al. Prevention and treatment of dementia or Alzheimer's disease by statins: a meta-analysis. *Dement Geriatr Cogn Disord*, 2007, 23:194-201.

7. DEPRESSION

7.1 A history of depression is associated with increased dementia risk

People with depression or high depressive symptoms have, on average, a higher risk of developing dementia. Evidence suggests that treatment with antidepressants improves cognition in people with depression, but it is not known if treatment prevents dementia.

This section describes the current evidence that depression increases dementia risk and provides links to resources for health and lifestyle professionals and for clients.

7.2 Depression and dementia – the evidence

A history of depression has been associated with an increased risk of Alzheimer's disease and dementia in several studies [1,2]. Some studies, however, have failed to replicate this finding.

In a study of 1,953 people with Alzheimer's disease, depressive symptoms before dementia onset were twice as common as in controls [3]. It may be that early awareness of cognitive decline contributes to depression, but it seems that this does not fully explain the association. A review of all epidemiological studies also reported an association between depression and Alzheimer's disease even when those episodes of depression had been more than 10 years before the onset of Alzheimer's disease [4].

It is well known that people with depression, especially older adults, have reduced cognitive performance and that many people with dementia also have depression. It is therefore difficult to determine whether depression is a risk factor for dementia or whether it is a prodromal symptom [1,2].

Depression is associated with elevated cortisol production, which may directly damage the hippocampus and increase the risk of dementia. Conversely, depression may arise secondary to dementia due to frontostriatal damage from cerebrovascular or neurodegenerative pathology [1,2]. Recent studies have also suggested that people with depression have enhanced deposition of beta amyloid plaques [1], and depression has been associated with vascular changes in the brain, which may mediate the association [5].

The treatment of depression seems to improve cognitive function, but it may not return cognition to normal levels even when the depression is in remission [1,2]. Whether the treatment of depression decreases the risk of dementia among people with depressive symptoms has not yet been studied, but it is clearly important to identify and treat depression. Preventing new episodes of depression may be a useful preventative approach to dementia [6].

7.3 Depression resources for professionals

Depression is a common disorder in the Australian community. The best outcomes are likely when adequate treatment is provided over a long enough time. Potentially reducing the risk of dementia is only one of the many reasons to identify and effectively treat depression. Clients at risk should be referred to their doctor.

The [Black Dog Institute](#) health professionals website has a wide range of information and resources, including information, education and a newsletter.

The [beyondblue](#) website has a wide range of information on depression for health professionals, including fact sheets and resources, information about training opportunities, assessment scales, journal articles and reports.

The Royal Australian and New Zealand College of Psychiatrists provide [clinical practice guidelines for the treatment of depression](#), which include detailed treatment recommendations. The [RANZCP](#) website also has clinical guidelines for other related mental disorders.

The [beyondblue Fact sheet 24 - Help for depression under Medicare](#) provides an overview of the *Better Access to Psychiatrists, Psychologists and GPs through the MBS' initiative (commonly referred to as Better Access to Mental Health Care)* focusing on who is eligible, what services are involved and what the cost is.

7.4 Depression resources for clients

[Coping with depression. Australian treatment guide for consumers and carers](#) is an information booklet from the RANZCP that provides information about depression, the treatments available and where to go for more information.

[What is a depressive disorder?](#) is an information booklet from the Australian Government Department of Health and Ageing, explaining what depression is, what causes it, how it is treated and where to go for help.

The [Black Dog Institute](#) public website has a wide range of information and resources on depression, and an online education program for consumers.

The [beyondblue](#) website has a wide range of information on depression for consumers.

The [Reach Out Australia](#) website has a range of information and resources for young people experiencing mental health issues.

7.5 References

1. Middleton LE, Yaffe K. Promising strategies for the prevention of dementia. *Arch Neurol*, 2009, 66:1210-1215.
2. Hughes TF, Ganguli M. Modifiable midlife risk factors for late-life cognitive impairment and dementia. *Curr Psychiatry Rev*, 2009, 5:73-92.
3. Green R, et al. Depression as a risk factor for Alzheimer's disease. The MIRAGE study. *Arch Neurol*, 2003, 60:753-759.
4. Jorm A, et al. Psychiatric history and related exposures as risk factors for Alzheimer's disease: a collaborative re-analysis of case-control studies. EURODEM risk factors research group. *Int J Epidemiol*, 1991, 20:S43-S47.
5. Woodward M, et al. *Dementia risk reduction: the evidence*. 2007, Alzheimer's Australia, Canberra.
6. Scalco MZ, van Reekum R. Prevention of Alzheimer disease. Encouraging evidence. *Can Fam Physician*, 2006, 52:200-207.

8. DIABETES

8.1 Diabetes increases dementia risk

Diabetes is associated with an increased risk of developing dementia, including Alzheimer's disease. The ability of treatment of diabetes to reduce dementia risk has not been established.

This section describes the current evidence that suggests assessment and control of diabetes are important for reducing dementia risk and provides links to resources for health and lifestyle professionals and for clients.

8.2 Diabetes and dementia risk – the evidence

Diabetes, especially type 2, and pre-diabetes syndromes appear to be risk factors for dementia. A recent review concluded that type 2 diabetes is associated with increased risk of Alzheimer's disease and vascular dementia [1]. Another review reported meta-analyses that found diabetes was associated with a 47% increased risk of any dementia, a 39% increased risk of Alzheimer's disease, and a 138% increased risk of vascular dementia [2].

Diabetes has also been associated with greater risk for cognitive decline in several studies following people for 2 – 7 years [2]. Global cognitive function and executive function have been shown to be affected by diabetes.

Few studies have examined the effect of treatment of diabetes on dementia risk and the results are mixed [2]. The effect of the duration of diabetes on dementia risk has also not been determined, however duration has been associated with increased risk of cognitive decline [2].

Insulin resistance has also been associated with increased dementia risk. Insulin inhibits the degradation of beta amyloid, thus increasing the formation of plaques. Hyperinsulinemia may also activate inflammation in the brain, thereby increasing the risk of dementia [2].

The mechanisms underpinning the association between diabetes and dementia risk are unclear. Multiple processes may be involved, such as the effects of insulin resistance, vascular pathology, oxidative stress, glucocorticoids excess and inflammation [1,2].

8.3 Diabetes resources for professionals

Everyone, including those concerned about their risk of developing dementia, should be educated about lifestyle choices to help prevent diabetes and the importance of screening. Diabetes, if present, should be treated as appropriate. Diabetes increases the risk of dementia, as well as heart disease and a range of other conditions.

The [Diabetes Australia](#) website provides links to best-practice guidelines for the prevention, diagnosis and management of diabetes for health professionals.

The Department of Health and Ageing's [Prevention of Type 2 Diabetes Program](#) provides resources for assessing diabetes risk and referring patients to lifestyle modification programs.

8.4 Diabetes resources for patients

The [Mind your Health Checks](#) section of the Alzheimer's Australia Mind your Mind website provides information for the general public on diabetes and its relationship to dementia risk.

[Diabetes Australia](#) provides a range of information for consumers on diabetes.

The Better Health Channel provides fact sheets on [pre-diabetes](#) and [diabetes](#) for consumers.

8.5 References

1. Strachan MWJ, et al. The relationship between type 2 diabetes and dementia. *British Medical Bulletin*, 2008, 88: 131-146.
2. Lu F-P, et al. Diabetes and the risk of multi-system aging phenotypes: a systematic review and meta-analysis. *PLoS ONE*, 2009, 4(1): e4144. doi: 10.1371/journal.pone.0004144.

9. DIET

9.1 A brain healthy diet may reduce dementia risk

Diets low in saturated fat and high in antioxidant rich fruits and vegetables have been associated with better brain function and reduced risk of dementia. While the benefits of specific nutrients are not yet clear, what is good for the heart seems to be also good for the brain. A healthy diet is likely to contribute to brain health and may reduce the risk of developing dementia.

This section describes the current evidence that suggests a healthy diet is important for reducing dementia risk and provides links to resources on healthy eating promotion for health and lifestyle professionals and resources for clients.

9.2 Diet and dementia risk – the evidence

There is growing evidence that a healthy diet is important for brain health and may help reduce the risk of developing dementia. In particular, avoiding saturated fat and including unsaturated fats, fruits and vegetables and fish may be important for brain health.

Several prospective studies have found that high intakes of saturated and transunsaturated (hydrogenated) fats are associated with increased risk of dementia, while higher intake of polyunsaturated and monounsaturated fats is associated with reduced risk of dementia [1].

The omega 3 fatty acids contained in fish oils are thought to reduce inflammation in the brain and promote neurogenesis. Results are conflicting as to whether omega 3 protects against dementia [2], but several studies have shown an association between higher fish consumption and lower dementia risk.

Antioxidants may help protect against oxidative damage, considered part of the pathology of Alzheimer's disease. Lower levels of antioxidants have been observed in people with Alzheimer's in cross-sectional studies [3]. Prospective studies provide some evidence of lower dementia risk for higher intake of specific antioxidants, but results are conflicting [3].

Several studies have found a lower risk of dementia is associated with higher intake of fruit and vegetables and higher adherence to the Mediterranean diet [4].

Folate and vitamin B12 deficiencies have been associated with cognitive impairment and dementia. However, whether increasing intake or using supplements of folate or vitamin B12 reduces the risk of developing dementia is as yet unknown [5]. Deficiencies or elevated homocysteine (an amino acid linked to folate and vitamin B12 levels) should be treated.

There is insufficient evidence to promote a specific diet for reducing dementia risk. However, the Australian Dietary Guidelines recommend avoiding saturated fat and eating plenty of fruits and vegetables [6]. This general healthy diet is likely to help reduce the risk of dementia, in addition to cardiovascular disease and other health conditions.

9.3 Promoting healthy diets in professional practice

Everyone, including those concerned about their dementia risk, should be encouraged to eat healthily, minimise saturated fat intake and increase fruit and vegetable consumption.

Clients should be advised to follow the [Dietary Guidelines for Australians](#). These recommend that all adults:

- ◆ Eat plenty of vegetables, legumes and fruits
- ◆ Eat plenty of cereals, preferably wholegrain
- ◆ Include lean meat, fish, poultry and/or alternatives
- ◆ Include reduced-fat milks, yoghurts, cheeses and/or alternatives.
- ◆ Drink plenty of water
- ◆ Limit saturated fat and moderate total fat intake
- ◆ Choose foods low in salt
- ◆ Consume only moderate amounts of sugars and foods containing added sugars

Referral to a dietician may be appropriate for some people. An Accredited Practising Dietician in your area can be found on the [Dieticians Association of Australia](#) website, which also provides information on healthy eating.

The [SNAP Guide](#) outlines the recommendations for assessment and intervention to help patients improve their diet. The SNAP guide was developed for general practitioners, but much of the information could be used by other professionals.

The [Australian Government Department of Health and Ageing](#) website provides a range of information and resources related to nutrition, and links to state and territory health department websites and other nutrition related organisations and services.

9.4 Healthy diet resources for clients

The [Mind your Diet](#) section of the Alzheimer's Australia Mind your Mind website provides information for the general public on healthy eating and its relevance to dementia risk.

The [Australian Dietary Guidelines](#) highlight the groups of foods and lifestyle patterns that promote good nutrition and health. Booklets and pamphlets summarising the guidelines for the general public are available at the website.

The [Heart Foundation's Healthy Eating](#) website provides information, advice and recipes to help patients/clients enjoy a healthy diet.

The [Nutrition Australia](#) website provides information about food and nutrition, related services and activities, and healthy recipes.

The [Go for 2&5 Campaign](#) website provides tips and recipes for increasing fruit and vegetable consumption.

Alzheimer's Australia has an Update Sheet [What you eat and drink and your brain](#) that provides information about the role of diet in potentially reducing the risk of developing dementia and in improving the health of people with dementia.

9.5 References

1. Solfrizzi V, et al. Dietary fatty acids intake: possible role in cognitive decline and dementia. *Experimental Gerontology*, 2005, 40:257-270.
2. Issa AM. The efficacy of omega–3 fatty acids on cognitive function in aging and dementia: a systematic review. *Dement Geriatr Cogn Disord*, 2006, 21:88-96.
3. Scalco MZ & van Reekum R. Prevention of Alzheimer disease. Encouraging evidence. *Can Fam Physician*, 2006, 52:200-207.
4. Lee Y, et al. Systematic review of health behavioral risks and cognitive health in older adults. *Int Psychogeriatr*, 2009, doi: 10.1017/S1041610209991189.
5. Woodward M, et al. *Dementia risk reduction: the evidence*. 2007, Alzheimer's Australia, Canberra.
6. National Health and Medical Research Council. *Dietary guidelines for Australian adults*. 2003, Commonwealth of Australia, Canberra.

10. HEAD INJURY

10.1 Head injury increases the risk of dementia

Head injury, with loss of consciousness, is associated with an increased risk of subsequent cognitive decline and dementia.

This section describes the current evidence that suggests preventing head injury increases dementia risk and provides links to further information.

10.2 Head injury and dementia – the evidence

Several studies have examined the association between head injury and risk of dementia in late-life. Head injury has been shown to increase the risk of cognitive decline, dementia and Alzheimer's disease [1]. The association may be mediated by the severity of head injury, whether loss of consciousness occurs, the time between head injury and dementia onset and genetic factors [1]. Boxing has also been associated with head injury and a form of dementia (dementia pugilistica), but not with Alzheimer's disease [2].

Long term follow up of US World War II veterans who were admitted to hospital during service showed that those who had suffered a head injury with loss of consciousness had an increased risk of subsequently developing Alzheimer's disease compared to those admitted for other reasons. Those with moderate head injuries had 2.3 times the risk and those with severe head injuries had 4.5 times the risk [3].

A meta-analysis of case control studies found a 58% increased risk of Alzheimer's disease for those with a history of head injury [4]. The association was found to be significant only in males. Males with a history of head injury had 2.3 times the risk of Alzheimer's, compared to 0.9 times the risk for females with a history of head injury. A possible explanation for the gender difference is the role of female hormones, which may confer a neuroprotective and neuroregenerative effect [4].

Several mechanisms have been proposed to explain the association between head injury and increased dementia risk. Damage to the blood brain barrier, increased oxidative stress, neuronal loss, increased enzyme activity leading to increased beta-amyloid deposition and increased tau pathology have all been implicated [1,2].

10.3 Prevention of head injury in professional practice

Prevention of head trauma is important for a number of reasons, including its potential effect on dementia risk. Children and adults should be encouraged to follow current safety advice and legislation to prevent head injury. This includes wearing helmets when riding bicycles or motorcycles, or participating in contact sports or other activities associated with a high risk of falls, and preventing falls in older adults.

Older people are more likely to suffer head or other injury due to falls, and the risk factors include:

- ◆ Stroke
- ◆ Parkinson's disease
- ◆ Arthritis
- ◆ Postural hypotension
- ◆ Dizziness
- ◆ Diabetes
- ◆ Depression
- ◆ Dementia
- ◆ Incontinence

- ◆ Poor nutrition
- ◆ Poor balance
- ◆ Weak muscles
- ◆ Poor eyesight
- ◆ Some medications and polypharmacy
- ◆ Foot problems and inadequate footwear

Health and lifestyle professionals can assist clients to prevent falls by:

- ◆ providing advice about exercise, particularly balance and strength exercises
- ◆ referring clients to a podiatrist, ophthalmologist, occupational therapist, physiotherapist, or other appropriate health professional
- ◆ referring clients to their local council, Commonwealth Carelink, Independent Living Centre or other appropriate agency for advice about services and supports available to them

10.4 Injury prevention information for clients

The [Mind your Head](#) section of the Alzheimer's Australia Mind your Mind website provides information for the general public on the links between head injury and dementia risk and tips for preventing head injury.

The Better Health Channel [Accidents and Injury – Reducing the Risks](#) web page provides information for consumers about preventing injury in the car, as a pedestrian, as a cyclist, in the workplace, at home, and when participating in sport and leisure activities.

The Better Health Channel [Falls Prevention for Older People](#) web page provides information for consumers about preventing falls through appropriate exercise, improving safety inside and outside the home, and managing health conditions and medications that may increase falls risk.

[Don't fall for it. Falls can be prevented! A guide to preventing falls for older people](#) is a booklet from the Australian Government Department of Health and Ageing. The booklet is aimed at community dwelling older people and their families and carers. It contains information about reducing falls risk factors relating to a person and their health and falls risk factors relating to the environment. It also describes what to do in the event of a fall.

The National Prescribing Service provides information for consumers about [avoiding falls caused by the side effects of medications](#).

10.5 References

1. Hughes TF, Ganguli M. Modifiable midlife risk factors for late-life cognitive impairment and dementia. *Curr Psychiatry Rev*, 2009, 5:73-92.
2. Woodward M, et al. *Dementia risk reduction: the evidence*. 2007, Alzheimer's Australia, Canberra.
3. Plassman BL, et al. Documented head injury in early adulthood and risk of Alzheimer's disease and other dementias. *Neurology*, 2000, 55:1158-1166.
4. Fleminger S, et al. Head injury as a risk factor for Alzheimer's disease: the evidence 10 years on; a partial replication. *J Neurol Neurosurg Psychiatry*, 2003, 74:857-862.

11. MENTAL STIMULATION

11.1 Mental stimulation may reduce dementia risk

Higher participation in mentally stimulating activities is associated with better cognitive function and reduced risk of cognitive decline and dementia. Those with a history of higher education, mentally demanding occupations or participation in mentally challenging leisure activities are consistently found to have a lower risk of developing dementia. Mental activity is likely to contribute to brain reserve, providing increased capacity to maintain normal cognitive function in the face of neurodegenerative disease. This may reduce the risk or delay the onset of dementia.

This section describes the current evidence that suggests mental activity is important for reducing dementia risk and provides links to resources for health and lifestyle professionals and resources for clients.

11.2 Mental activity and dementia risk – the evidence

There is consistent evidence that higher participation in mentally stimulating activities may protect against cognitive decline and dementia in old age.

A reduced risk of dementia is associated with:

- ◆ Higher levels of education
- ◆ Mentally demanding occupations
- ◆ More cognitively stimulating leisure activities
- ◆ Higher intelligence

Meta-analysis suggests around half the risk of dementia for those with high educational attainment compared to low, those with high occupational status compared to low, and those participating in more cognitively stimulating leisure activities compared to those participating in few such activities [1].

Importantly for older or retired people, increased complex mental activity in late life is associated with lower dementia risk independent of other predictors such as education [1]. A dose–response relationship is also evident, so the more activity that can be done, the better.

Complex mental activity across the lifespan may reduce the risk or delay the onset of dementia by improving brain reserve [1,2]. Mental activity, challenge and learning stimulate neurogenesis and new synapses [2]. Complex cognitive activity contributes to neurological brain reserve (increased synapses, neural numbers and brain volume) and behavioural brain reserve (flexible cognitive strategies). This reserve may allow normal cognitive function to continue for longer in the face of underlying neural dysfunction or degeneration [3].

Commercial brain training games and programs have as yet not been shown to reduce the risk of dementia [4].

Activities that combine mental, social and physical components may provide even greater protection against dementia [2].

11.3 Promoting mental activity in professional practice

People concerned about their risk of developing dementia should be encouraged to remain mentally active throughout life. Older clients or those who are retired may be at particular risk of becoming less mentally active and should be encouraged to maintain regular participation in cognitively challenging activities.

Mental activity should be regular, reasonably complex and varied. It should involve continued new learning. It should also be something the person enjoys so it can be sustained. It could include anything that challenges the mind, for example:

- ◆ Working (full-time or part-time) or volunteering
- ◆ Hobbies – gardening, art, craft, collecting, etc
- ◆ Taking a course, learning a language or a new skill

Clients wanting to increase their cognitive activities could be referred to their local council or adult education provider. Health and lifestyle professionals can also provide follow up to monitor progress and encourage continued participation.

The 5As model based on interventions shown to be effective in motivating behaviour change could be applied to mental activity.

- ◆ Ask about the person's current mentally challenging activities
- ◆ Assess their level of risk and their readiness to change
- ◆ Advise them of the need for mental stimulation and provide information about what they can do
- ◆ Assist them to find activities they will enjoy and to plan how they will increase their level of mental activity
- ◆ Arrange referral to any appropriate services and follow up appointments

The 5As model for detection, assessment and management of risk factors is outlined in the [SNAP guide](#).

11.4 Mental activity resources for clients

The [Mind your Brain](#) section of the Alzheimer's Australia Mind your Mind website has information for the general public explaining why we believe mental activity is important and provides ideas and resources for brain exercise.

Your local council should be able to advise your clients about providers of locally available courses, activities, groups, volunteer opportunities, etc.

The internet is a great source of brain games. For example:

- ◆ The [Lipton BrainTrain](#) website has puzzles and games to challenge memory, maths skills and reaction time. Patients can play just for fun or sign up to play against others.
- ◆ The [SBS ADbc quiz](#), based on the TV program, tests history knowledge.
- ◆ The [Games for Brain](#) website has many different types of games challenging different cognitive skills.
- ◆ The [BrainBashers](#) website has games, puzzles, jigsaws, brain teasers riddles, optical illusions and more.

Brain games and puzzles can also be found in daily newspapers and in various puzzle books available at newsagents and bookstores.

Alzheimer's Australia has an Update Sheet [Keep Your Brain Active: Mental Exercise and Dementia](#) that provides information about the role of mental activity in reducing the risk of developing dementia and in improving the quality of life of people with dementia.

U3A - the University of the Third Age - is an international movement that encourages retired people to take part in lifelong learning. They offer many educational, creative and leisure activities to their members to encourage positive ageing. The [University of the Third Age Online](#) website provides up-to-date contacts for all Australian and New Zealand U3As. U3A Online also offers short online courses and many other resources for older people, especially those who are geographically, physically or socially isolated.

The [Sharp Brains](#) website has independent information about the many 'brain training' products commercially available. The University of New South Wales has an [information sheet](#) summarising what the research to date says about the effectiveness of these products.

Dr Michael Valenzuela's book, [It's Never Too Late to Change Your Mind. The Latest Medical Thinking on What You Can Do to Avoid Dementia](#), published by ABC Books, explains dementia and the major risk factors and provides practical advice for consumers on keeping active.

11.5 References

1. Valenzuela MJ & Sachdev P. Brain reserve and dementia: a systematic review. *Psychological Medicine*, 2006, 36: 441-454.
2. Valenzuela MJ. *It's never too late to change your mind*. 2009, ABC Books, Sydney.
3. Welsh-Bohmer KA & White CL. Alzheimer disease: What changes in the brain cause dementia? *Neurology*, 2009, 72: e21-e23.
4. Valenzuela MJ & Sachdev P. Can cognitive exercise prevent the onset of dementia? Systematic review of randomized clinical trials with longitudinal follow-up. *Am J Geriatr Psychiatry*, 2009, 17: 179-187.

12. PHYSICAL ACTIVITY

12.1 Physical exercise may reduce dementia risk

Regular physical exercise is associated with better brain function and reduced risk of cognitive decline and dementia. This has been shown in many studies for exercise undertaken in midlife and in late-life. Even simple exercise like walking has been shown to be beneficial. Getting the recommended 30 minutes of physical activity on most days is likely to contribute to brain health and may reduce the risk of developing dementia.

This section describes the current evidence that suggests physical activity is important for reducing dementia risk and provides links to resources on physical activity promotion for health and lifestyle professionals and resources for clients.

12.2 Physical Activity and dementia risk – the evidence

There is growing evidence that physical activity may protect against cognitive decline and dementia in old age.

Several longitudinal studies have found that older people who exercise are less likely to experience cognitive decline or develop dementia. For example, exercising at least 3 times per week in people over age 65 was associated with a 38% reduced risk of developing dementia after 6 years follow up [1]. In a randomised controlled trial of people aged 50 and over with subjective memory impairment, a 6 month program of physical activity resulted in improved cognition at the end of the trial and after 18 months follow up [2].

Regular exercise during midlife may also protect against later developing dementia. In one study, exercising at least twice a week at midlife was associated with a 52% reduced risk of dementia at age 65-79 [3].

Several mechanisms have been proposed to explain the association between physical inactivity and cognitive decline [4,5].

- ◆ Inactivity increases the risk of vascular disease and vascular risk factors. Vascular disease, in turn, increases the risk and severity of cognitive decline and dementia.
- ◆ Inactivity is associated with higher levels of inflammatory markers in the blood (e.g. C-reactive protein). Inflammation has been associated with an increased risk of cognitive decline and dementia.
- ◆ Physical activity appears to stimulate neurogenesis, providing another pathway by which physical activity could protect against cognitive decline and dementia.

Regular physical exercise provides a range of health benefits. The evidence supporting the beneficial effects of physical activity on brain health provides yet more reasons to encourage regular physical activity among people of all ages.

12.3 Promoting physical activity in professional practice

Everyone, including those concerned about their risk of developing dementia, should be encouraged to be physically active. All adults should be advised to participate in 30 minutes of moderate activity on most, preferably all days of the week. More vigorous exercise may confer additional health and disease prevention benefits.

The [SNAP guide](#) provides recommendations and advice on implementing interventions to encourage patients to be more physically active. The SNAP guide was developed for general practitioners, but much of the information could be used by other professionals.

The [Australian Association for Exercise and Sports Science](#) website provides information on services provided by Exercise Physiologists, and allows you to locate an Exercise Physiologist in your area.

The [Australian Government Department of Health and Ageing](#) website provides a range of information and resources related to physical activity, and links to state and territory health department websites, which provide information about the many state based physical activity initiatives and resources.

12.4 Physical activity resources for clients

The [Mind your Body](#) section of the Alzheimer's Australia Mind your Mind website provides information for the general public on physical activity and its link to dementia risk, and provides links to some physical activity programs.

The [Heart Foundation](#) runs walking groups and exercise programs nationally, and the website provides consumer information on physical activity.

[An active way to better health: National Physical Activity Guidelines for Adults](#) from the Australian Government Department of Health and Ageing outlines the minimum levels of physical activity required to gain a health benefit and ways to incorporate physical activity into everyday life.

[Choose Health: Be Active: A physical activity guide for older Australians](#) is a booklet that offers ideas for keeping active at no cost, including balance and strength exercises that can be done at home.

Alzheimer's Australia has an Update Sheet [Keep on Moving: Physical Exercise and Dementia](#) that provides information about the role of physical activity in reducing the risk of developing dementia and in improving the health of people with dementia.

12.5 References

1. Larson EB, et al. Exercise is associated with reduced risk for incident dementia among persons 65 years of age and older. *Ann Intern Med*, 2006, 144:73-81.
2. Lautenschlager NT, et al. Effect of physical activity on cognitive function in older adults at risk for Alzheimer's disease. *JAMA*, 2008, 300:1027-1037.
3. Rovio S, et al. Leisure-time physical activity at midlife and the risk of dementia and Alzheimer's disease. *Lancet Neurol*, 2005, 4:705-711.
4. Jedrzewski MK, et al. Physical activity and cognitive health. *Alzheimers Dement*, 2007, 3:98-108.
5. Hendrie HC, et al. The NIH cognitive and emotional health project. Report of the critical evaluation study committee. *Alzheimers Dement*, 2006, 2:12-32.

13. SMOKING

13.1 Smoking increases dementia risk

Smoking is associated with an increased risk of dementia. Assisting smokers to quit may help reduce their risk of developing dementia.

This section describes the current evidence that suggests smoking increases dementia risk and provides links to resources on smoking cessation for health and lifestyle professionals and resources for clients.

13.2 Smoking and dementia risk – the evidence

Smoking has been demonstrated to be a risk factor for dementia. Two recent meta-analyses concluded that current smokers have an increased risk of any dementia, Alzheimer's disease, vascular dementia and cognitive decline compared to non-smokers [1,2].

Former smokers were not found to be at increased risk compared to never-smokers, suggesting quitting smoking may be beneficial for dementia risk [1,2].

A recent study examined the association between a history of heavy alcohol use and smoking and age of disease onset in 685 people with Alzheimer's disease. Heavy smoking (defined as one pack per day or more) was associated with a 2-3 year earlier onset of Alzheimer's disease [3].

The adverse health effects of exposure to secondhand smoke are similar to those of active smoking. There is some evidence that passive smoking may also be associated with increased risk of cognitive impairment and dementia [4].

These findings fit with other evidence of smoking as a risk factor for cardiovascular and cerebrovascular disease, stroke, increased oxidative stress, atherosclerosis and inflammation. Smoking may affect dementia risk via its negative effects on the cardiovascular system, increasing risk for both Alzheimer's disease and vascular dementia. Smoking may also interact with other cardiovascular risk factors in an additive manner [1,2].

The evidence suggests that reducing the risk of dementia is another good reason for smokers to be advised to quit.

13.3 Promoting smoking cessation in professional practice

People who smoke should be offered advice and assistance to quit. Smoking cessation is associated with reduced risk of cardiovascular disease and cancer, and may reduce the risk of developing dementia.

The [Smoking Cessation Guidelines for Australian General Practice](#) provide effective strategies to identify smokers and assist them to stop smoking.

The [SNAP guide](#) provides recommendations and advice on implementing these interventions and encouraging patients to quit smoking. The SNAP guide was developed for general practitioners, but much of the information could be used by other professionals.

Further information and resources are available from the [QUIT](#) website or Quitline on 13 7848 (13 QUIT).

13.4 Smoking cessation resources for clients

The [Mind your Habits](#) section of the Alzheimer's Australia Mind your Mind website provides information on the link between dementia and smoking and resources for consumers.

Quitline (13 7848 or 13 QUIT) is a confidential telephone service providing information, support and advice for quitting smoking. Trained advisors are available 8 am to 8 pm Monday to Friday.

The [National Tobacco Campaign](#) website provides information and resources to help patients quit smoking.

The [Quit Victoria](#) website provides access to an online Quit Coach, a 4 step guide to quitting, information about attending Quit courses, and a free Quit Pack that can be ordered.

The [Quit SA](#) website provides resources to help people quit smoking, including a free Quit Pack, and information for schools, health professionals and workplaces on what they can do to help reduce smoking in the community.

The [Quit Tasmania](#) website provides resources for quitting, information on programs and campaigns and resources for young people.

13.5 References

1. Anstey KJ, et al. Smoking as a risk factor for dementia and cognitive decline: a meta-analysis of prospective studies. *American Journal of Epidemiology*, 2007, 166:367-378.
2. Peters R, et al. Smoking, dementia and cognitive decline in the elderly, a systematic review. *BMC Geriatrics*, 2008, 8:36 doi:10.1186/1471-2318-8-36.
3. Harwood DG, et al. The effect of alcohol and tobacco consumption, and apolipoprotein E genotype, on the age of onset in Alzheimer's disease. *Int J Geriatr Psychiatry*, 2009, doi: 10.1002/gps.2372.
4. Llewellyn DJ, et al. Exposure to secondhand smoke and cognitive impairment in non-smokers: national cross sectional study with cotinine measurement. *BMJ*, 2009, 338: b462. doi: 10.1136/bmj.b462.
5. Zwar N, et al. *Smoking cessation guidelines for Australian general practice*. 2004, Australian Government Department of Health and Ageing, Canberra.

14. SOCIAL ACTIVITY

14.1 Social interaction may reduce dementia risk

Being more socially active is associated with reduced risk of cognitive decline and dementia. This has been shown in many studies measuring social engagement in different ways. Combining social activity with mental and/or physical activity may provide even greater benefit in reducing the risk of developing dementia.

This section describes the current evidence that suggests social activity is important for reducing dementia risk and provides links to resources for health and lifestyle professionals and for clients.

14.2 Social Activity and dementia risk – the evidence

A history of being more socially active is associated with a reduced risk of developing dementia. Participating in more social activities, having larger networks of friends and feeling less lonely have all been associated with a lower risk of developing dementia.

The association between social engagement and dementia risk has been shown by several studies, for example:

- ◆ Participation in high numbers of different leisure activities was associated with a 38% lower risk of developing dementia in 1,772 people over age 65 [1]. Activities included going to clubs, visiting friends or being visited, playing cards and community or volunteer work.
- ◆ Larger social networks were associated with a 36% lower risk of dementia in 2,249 elderly women [2].
- ◆ Loneliness (perceived isolation) was associated with more than double the risk of developing dementia in 823 older persons [3]. Loneliness was also associated with worse cognition at baseline and more rapid cognitive decline during follow-up.

Research suggests that social activities that also involve mental stimulation and/or physical activity can provide even greater benefit.

A study in Stockholm investigated mental, physical and social components of the leisure activities of people aged 75 years and older [4]. It found that all 3 components were associated with a reduced risk of dementia after 6 years follow up. Combining components offered the greatest benefit, with those whose activities included higher levels of 2 or all 3 components having a 47% reduced risk of dementia.

14.3 How does social activity reduce dementia risk?

Social activity may contribute to brain reserve. Interacting with other people involves many cognitive functions. As with other mentally stimulating activity, this is believed to help build up a reserve of healthy neurons and synapses that may protect against dementia.

14.4 Promoting social activity in professional practice

Everyone, including those concerned about their risk of developing dementia, should be encouraged to be socially active as far as they feel comfortable. Social isolation can increase the risk of dementia, heart disease and mental illness.

There are many reasons why a person may not be socially engaged. For example, patients who are older, geographically isolated, disabled or suffering chronic illness may be at higher risk of social isolation.

The 5As model for detection, assessment and management of risk factors, as outlined in the [SNAP guide](#), could be applied to social activity.

- ◆ Ask about the person's current social activities
- ◆ Assess their level of risk and their readiness to change
- ◆ Advise them of the need for social engagement and provide information about what they can do
- ◆ Assist them to find activities they will enjoy and to plan how they will increase their level of social activity
- ◆ Arrange referral to any appropriate services and follow up appointments

14.5 Social activity resources for clients

Your local council can provide information about clubs, groups, courses, etc in your local area that clients could get involved in.

The [Mind your Social Life](#) section of the Alzheimer's Australia Mind your Mind website has information for the general public explaining why we believe social activity is important and provides ideas for social engagement.

Alzheimer's Australia's booklet [Mind your Mind: How to keep your brain healthy and reduce your risk of dementia](#) lists a variety of tips for staying socially active.

The [Heart Foundation](#) runs walking groups and exercise programs nationally, that could help patients increase their social and physical activity.

14.6 References

1. Scarmeas N, et al. Influence of leisure activity on the incidence of Alzheimer's disease. *Neurology*, 2001, 57:2236-2242.
2. Crooks VC, et al. Social network, cognitive function, and dementia incidence among elderly women. *American Journal of Public Health*, 2008, 98:1221-1227.
3. Wilson RS, et al. Loneliness and risk of Alzheimer disease. *Archives of General Psychiatry*, 2007, 64:234-240.
4. Karp A, et al. Mental, physical and social components in leisure activities equally contribute to decrease dementia risk. *Dement Geriatr Cogn Disord*, 2006, 21:65-73.

APPENDIX

Dementia risk reduction summary sheet (one page) for distribution to clients



**MIND
YOUR
MIND**



**Alzheimer's
Australia**
Living with dementia

Mind your Mind to keep your brain healthy

We now have good evidence from research that shows adopting a 'brain healthy' lifestyle may reduce the risk of developing dementia. Alzheimer's Australia's **Mind your Mind**[®] program involves seven signposts to help keep your brain healthy and reduce your risk of developing dementia.

For more information on Alzheimer's Australia's **Mind your Mind** program contact us.

www.alzheimers.org.au

National Dementia Helpline 1800 100 500
An Australian Government initiative

There are a number of factors that increase your risk of developing dementia, including age and genes. Following the **Mind your Mind** signposts, can't guarantee that you won't get dementia, but it may help reduce your risk or delay the onset of dementia.

© Lynton Chubb Photography
August 2008 © Alzheimer's Australia Inc. ABN 796 2568 2771 A.RBN 066 779 557

Mind your Mind[®] signposts



Mind your Brain

Challenge your brain every day with new things

- ◆ Play crosswords, number and word puzzles or board games
- ◆ Learn a language, a musical instrument or enrol in a course
- ◆ Experience new activities like going to plays, concerts, museums and galleries



Mind your Diet

Enjoy healthy eating by choosing:

- ◆ A wide variety of foods
- ◆ Vegetables, fruits, legumes (dried peas, beans, chickpeas and lentils), wholegrain breads and cereals
- ◆ Lean meats, chicken, fish and reduced-fat dairy products
- ◆ Oils such as olive, sunflower, safflower, soy, canola and flaxseed



Mind your Body

Be active every day in ways you find enjoyable

- ◆ Walk or cycle to work or around your neighbourhood
- ◆ Go dancing, jogging or swimming
- ◆ Join a gym, yoga, tai chi or pilates group
- ◆ Do some gardening
- ◆ Play sport socially or competitively



Mind your Health Checks

**Know your blood pressure, blood cholesterol,
blood sugar and weight**

- ◆ These can all increase your risk of developing dementia if they are too high. Ask your doctor to check these and help you manage any problems.



Mind your Social Life

Participate in social activities, stay socially connected

- ◆ Catch up with family and friends
- ◆ Join a club or group
- ◆ Participate in community events or become a volunteer



Mind your Habits

Avoid 'bad habits'

- ◆ Don't smoke
- ◆ If you drink alcohol, drink in moderation



Mind your Head

Protect your head from serious injury

- ◆ Take care as a pedestrian and always wear a seatbelt
- ◆ Use protective headgear when riding, skating and playing sport



An Australian Government Initiative

Visit the Mind your Mind website at
www.mindyourmind.org.au
for further information about dementia risk reduction

Visit the Alzheimer's Australia website at
www.alzheimers.org.au
for comprehensive information about

- dementia and care
- information, education and training
- other services offered by member organisations

Or for information and advice contact the
National Dementia Helpline on 1800 100 500

Visit the Dementia Collaborative Research Centres website at
www.dementia.unsw.edu.au
for further information about the people involved and the research activities